



Immunology immunoglobulin referral form

Infusion pharmacy
Phone: 1-877-342-9352
Fax: 1-888-594-4884

✂ Please detach before submitting to a pharmacy—tear here.

PATIENT INFORMATION			
IG specialist: Name:		Phone:	
Patient: see attached	Gender: Male Female		
Patient name:		DOB:	SSN:
Address:		City:	
State:	ZIP:	Phone:	Cell:
Emergency contact:		Phone:	Relationship:
Insurance: Front and back of insurance card to follow			
Primary Insurance:	Phone:	Policy #:	Group:
Secondary Insurance:	Phone:	Policy #:	Group:
Primary diagnosis: Combined immunodeficiency Common hypogammaglobulinemia Common variable immunodeficiency Immune thrombocytopenic purpura Immunodeficiency with increased IgM Wiskott-Aldrich syndrome Other:			
Medical assessment: Height:		Weight:	lbs kg
Current medications? Yes No If yes, list or attach:			
Allergies:			

PRESCRIPTION ORDERS Medication infused per PI recommended rate and via rate controlled device per therapy			
Immune Globulin: No preference Preferred product:			
Directions: Infuse IV Infuse SC <input checked="" type="checkbox"/> Per manufacturer guidelines or as written:		<input checked="" type="checkbox"/> May round to the nearest 5gm vial size	
Initial:	gm/kg divided over	days; OR	Other:
Ongoing:	gm/kg divided over	days, every	weeks for cycles; OR Other:
Quantity/Refills: 1-month supply; refill x 12 months unless otherwise noted Other:			
Pre-medications 30 minutes before start of IG:			
Acetaminophen PO	325 mg 650 mg	mg	DiphenhydrAMINE PO 25mg 50mg mg
Hydration, solution:	Volume:	mL/hr:	Other:
Nursing and other orders:			
<input checked="" type="checkbox"/> Administer IVIG or teach SCIG self-administration, via pump <input checked="" type="checkbox"/> Ambulatory pump if required for infusion			
<input checked="" type="checkbox"/> Initiate access device (<i>insert peripheral IV, SC needles, access implanted port, or use existing PICC</i>)			
<input checked="" type="checkbox"/> Flush IV with 5mL 0.9% NaCl; Heparin 100 units/mL OR 10 unit/mL, 3-5mL PRN			
Obtain labs (list): Lab frequency: Once Monthly Other:			
<input checked="" type="checkbox"/> Anaphylaxis kit order Infusion Reaction Management x1 year			
Mild	<ul style="list-style-type: none"> • Slow infusion rate by 50% until symptoms resolve. Resume at previous rate as tolerated. <input checked="" type="checkbox"/> DiphenhydrAMINE PO 25mg 50mg mg Dispense diphenhydrAMINE 25mg capsules x 4 		
Moderate	<ul style="list-style-type: none"> • Stop Infusion, resume at 50% rate when symptoms resolve <input checked="" type="checkbox"/> DiphenhydrAMINE IV 25mg 50mg mg Dispense diphenhydrAMINE 50mg vial x 1 		
Severe (Anaphylaxis) *Call 911* Notify prescribing physician	<ul style="list-style-type: none"> • Stop infusion and remove tubing from access device to prevent further administration • Initiate 0.9% NaCl 500mL/hr IV OR mL/hr <input checked="" type="checkbox"/> Dispense 0.9% NaCl 500mL x 1 • Administer EPINEPHrine 1mg/mL by weight (Wt.) as an IM injection into the lateral thigh Wt > 66lbs (30kg) Wt 33 to 66 lbs (15 to 30kg) Wt < 33lbs (15kg) 0.3mg/0.3mL 0.15mg/0.15mL 0.01mg/kg • Repeat EPINEPHrine in 5 to 15 minutes if symptoms persist • Administer CPR if needed until EMS arrives <input checked="" type="checkbox"/> Dispense EPINEPHrine x 2: 1mg vial Pen JR 0.15mg Pen 0.3mg Other: 		

PHYSICIAN INFORMATION			
Name:		Practice:	
Address:		City:	State: ZIP:
Phone:	Fax:	NPI:	Contact:
By signing, I certify/recertify that the above therapy, products and services are medically necessary and that this patient is under my care. I have received authorization to release the above referenced information and medical and/or patient information relating to this therapy. Pharmacy has my permission to contact the insurance company on my behalf to obtain authorization for patient.			
Substitution permissible Signature:		Date:	Dispense as written Signature:

This form is not a valid prescription in Arizona.